




Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Journal Pre-proof



Holistic care for patients with severe coronavirus disease 2019: An expert consensus

Nursing Department of Tongji Hospital Affiliated to Tongji Medical College Hust,
Nursing Department of Peking Union Medical College Hospital, Intensive Care
Professional Committee of the Chinese Nursing Association, Writing Committee
Members, Hui Wang, Tieying Zeng, Xinjuan Wu, Hong Sun

PII: S2352-0132(20)30048-X

DOI: <https://doi.org/10.1016/j.ijnss.2020.03.010>

Reference: IJNSS 484

To appear in: *International Journal of Nursing Sciences*

Received Date: 18 March 2020

Accepted Date: 31 March 2020

Please cite this article as: Nursing Department of Tongji Hospital Affiliated to Tongji Medical College Hust, Nursing Department of Peking Union Medical College Hospital, Intensive Care Professional Committee of the Chinese Nursing Association, Writing Committee Members, H. Wang, T. Zeng, X. Wu, H. Sun, Holistic care for patients with severe coronavirus disease 2019: An expert consensus, *International Journal of Nursing Sciences* (2020), doi: <https://doi.org/10.1016/j.ijnss.2020.03.010>.

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Copyright © 2020, Chinese Nursing Association. Production and hosting by Elsevier B.V. All rights reserved.

Dothead: **Responding to the COVID-19 Epidemic**

Title: **Holistic care for patients with severe coronavirus disease 2019: An expert consensus**

Nursing Department of Tongji Hospital Affiliated to Tongji Medical College Huat, Nursing Department of Peking Union Medical College Hospital, Intensive Care Professional Committee of the Chinese Nursing Association.
Writing Committee Members: Hui Wang^a, Tieying Zeng^a, Xinjuan Wu^{b*}, Hong Sun^b

^a*Nursing Department, Tongji Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China*

^b*Nursing Department, Peking Union Medical College Hospital, Chinese Academy of Medicine Sciences, Beijing, China*

☆ This article is based on of a previously published article in Chinese [Chinese Journal of Nursing, 2020, 55(3): 337-42. DOI: 10.3761/j.issn.0254-1769.2020.03.003].

***Corresponding author.**

E-mail address: wuxinjuan@sina.com(X.-J. Wu)

Ethical Consideration

The project was conducted in accordance with the Declaration of Helsinki (2013).

Acknowledgement

The authors would like to acknowledge the dedication of the nurses in fighting against COVID-2019.

Conflicts of interest

None.

Funding

None.

Holistic care for patients with severe coronavirus disease 2019: An expert consensus

【Abstract】

Objective To standardize the holistic care for patients with severe coronavirus disease 2019 (COVID-19).

Methods The consensus was preliminarily formed by combining relevant literature findings and frontline medical working experiences, and it was eventually confirmed by five rounds of online discussions and expert consultations.

Results This consensus included nursing assessment, nursing priorities, nursing goals, and thirteen key points of nursing procedures such as nursing of oxygen therapy and respiratory nursing.

Conclusion This scientific and practical consensus guideline can provide clinical guidance on the holistic nursing care of patients with severe COVID-19.

【Key words】 COVID-19; Expert Consensus; Holistic Nursing; Intensive Care

What is known?

The condition of patients with COVID-19 who need intensive care changes rapidly. These patients may lose their lives because infections often affect multiple system organs and lead to multiple organs dysfunction syndrome.

What is new?

More than twenty nursing experts in China discussed together and finally conformed a consensus on holistic nursing care of patients with severe COVID-19, which included nursing assessment, nursing priorities, nursing goals, and 13 key points of nursing such as nursing of oxygen therapy and respiratory nursing.

1. Introduction

At the end of 2019, an outbreak of coronavirus disease 2019 (COVID-19) was reported in Wuhan, China. As the epidemic continued to spread, it was listed by the World Health Organization (WHO) as a Public Health Emergency of International Concern [1], and China initiated a first-level response [2]. Due to the disease's high infectivity and pathogenicity and the high mortality rate of severely affected patients [3-4], nursing care of critically ill patients with COVID-19 is extremely difficult and requires high standards. Thus, The Chinese Government has proposed establishing a holistic nursing system for severe and critical patients to provide patient-centered care following modern nursing concepts, and utilizing nursing procedures as a fundamental framework and guide for clinical nursing care and nursing management. To standardize and guide holistic care of patients with COVID-19 in severe and critical condition and to

effectively preserve their physical and mental health, an expert consensus panel on Holistic Care of COVID-19 Patients in Severe and Critical Condition (hereafter “Consensus”) was jointly developed, led by the Chinese Nursing Association, and involving the Nursing Department of Tongji Hospital Affiliated with Tongji Medical College of Huazhong University of Science and Technology), the Nursing Department of the Peking Union Medical College Hospital of the Chinese Academy of Medical Sciences, and nursing experts dispatched by the Intensive Care Professional Committee of the Chinese Nursing Association to assist Wuhan.

2. Development of the Consensus

2.1. Target groups and protection levels

The Consensus applies to patients whose clinical condition is classified as either severe or critical according to the *Diagnosis and Treatment Protocol for Novel Coronavirus Pneumonia (Trial version 6)* [5]. When providing care to these patients, nurses should strictly implement personal protection measures. Level II protection should be adopted when working in isolation wards, and Level III protection measures should be adopted when performing procedures that may produce aerosols (e.g., sputum aspiration, tracheal intubation, noninvasive ventilation, tracheotomy, cardiopulmonary resuscitation, etc.).

2.2. Establishment of the Consensus

The Consensus was established jointly by 22 nursing experts, one medical expert, and one hospital infection control expert from different departments of 20 hospitals across China.

A draft of the consensus was developed after analysis, summary, and consolidation of information from a comprehensive review, which included the diagnosis and treatment protocols, management consensus, and related literature of COVID-19, as well as effective nursing measures and experiences gained by the experts during frontline care of patients in severe and critical condition.

Several revisions were made throughout five rounds of online discussion and expert consultations; a final version of the Consensus was then developed and reviewed by the medical expert and hospital infection control expert.

3. Contents of the Consensus

3.1. Key points in assessment

3.1.1. Medical history assessment

Assess the patient’s history of living pattern in the epidemic area, past medical history, treatment history, epidemiological history, and comorbidities, etc.

3.1.2. Physical assessment

- (1) Vital signs: Assess the patient's body temperature; blood pressure; pulse; blood oxygen saturation; breathing rate, rhythm, and depth; etc. [6].
- (2) Symptoms and signs: Assess the patient's level of consciousness, muscle pain, fatigue, cough, sputum, chest tightness, shortness of breath, diarrhea, etc.
- (3) Skin and mucosa: Assess the patient's skin color and elasticity, peripheral extremity circulation, presence of bleeding, etc.
- (4) Nutritional status: Assess the patient's food intake, presence of dehydration, etc.

3.1.3. Psychological assessment

Assess the patient's emotional response to the disease, cognitive changes, and compliance with protective measures.

3.1.4. Environmental assessment

Assess whether environmental factors can aggravate cross-infection, including inadequate bed spacing and the negative pressure ward, suboptimal protective measures for high-risk surgeries, and inadequate personal protective measures for all medical and nursing staff.

3.2. Health problems

3.2.1. Priority health problems

- (1) Impaired gas exchange – related to increased airway resistance and decreased lung compliance.
- (2) Hyperthermia – related to viral lung infection.
- (3) Activity intolerance – related to impaired lung function and imbalanced oxygen supply and consumption.
- (4) Potential complications – deep vein thrombosis, hyperglycemia, acute respiratory distress syndrome, septic shock, metabolic acidosis, coagulopathy, multiple organs dysfunction syndrome, stress ulcers, etc. [7].
- (5) Anxiety/fear – related to isolation and concerns over the disease's prognosis.

3.2.2. Other health problems

- (1) Ineffective airway clearance – related to excessive mucus and ineffective cough.
- (2) Diarrhea – related to intestinal dysfunction due to COVID-19 infection.
- (3) Impaired comfort – related to illness-related symptoms, such as chest tightness, gasping breaths and fatigue.
- (4) Imbalanced nutrition: less than body requirements – related to fever, diarrhea, reduced intake, etc.
- (5) Risk for impaired skin integrity – related to fever, humidity, poor peripheral circulation, and forced position.

- (6) Knowledge deficit – patients may have insufficient knowledge related to isolation concept, personal protection and COVID-19.

3.3. Nursing goals

- (1) Alleviating dyspnea and hypoxemia.
- (2) Ensuring effective coughing, with breathing gradually becoming stable and lung sounds clear.
- (3) Controlling body temperature effectively.
- (4) Relieving diarrhea.
- (5) Reducing chance of developing pressure injury during hospitalization.
- (6) Detecting and handling complications effectively.
- (7) Ensuring patient to have a stable emotion and gaining effective cooperation from patients when implementing therapeutic nursing interventions and rehabilitation programs.
- (8) Ensuring a gradual return to normal diet and maintaining a desired body weight at a standard level.
- (9) Maintaining patient comfort during hospitalization and satisfying their needs timely.
- (10) Improving patients' activity endurance.
- (11) Ensuring high compliance to isolation and effective self-protection among patients.

3.4. Nursing interventions

3.4.1. Providing optimal oxygen therapy and respiratory care

- (1) Promptly evaluate whether patients' respiratory distress and/or hypoxemia are relieved and provide patients with individualized respiratory support per medical orders based on their condition and tolerance [8].
- (2) Patients receiving nasal catheter oxygen or nasal high-flow oxygen therapy can wear a surgical or face mask to reduce the spread of viral aerosols [9]. Oxygen supply cannot be interrupted arbitrarily during nasal high-flow oxygen therapy [10]. Employ lung protective approaches during invasive mechanical ventilation, and apply a small tidal volume (4–8 ml/kg of ideal weight) and low inspiratory pressure (plateau pressure < 30 cmH₂O [1 cmH₂O = 0.098 kPa]) during the procedure to reduce ventilator-induced lung injury [5].
- (3) During administration of oxygen therapy, closely monitor patients' consciousness, heart rate, breathing characteristics (rhythm, rate, frequency, depth, synchronization of spontaneous breathing and ventilator), improvement of cyanosis, and oxygen therapy complications. Carefully monitor blood oxygen saturation or arterial blood gas analysis results [11] and promptly adjust oxygen flow rate, oxygen supply mode, and ventilator parameters accordingly, as per medical orders.
- (4) Ventilator and catheter care: (i) Passively humidify invasive ventilators using a heat-moisture-exchanger (HME) with a bacterial filter to reduce condensation in the circuit [12].(ii) Use of disposable ventilator circuits and exhalation valves is recommended, and installation of virus filters at both the inhalation and exhalation ends of the ventilator. Replace these filters only upon contamination or mechanical failure, as opposed to on a routine basis [12]. (iii) Do not disconnect the

ventilator. If the ventilator must be disconnected, for reasons such as replacing the sealed sputum-aspirating tube, it must first be set to standby mode. (iv) Use 75% alcohol to wipe and disinfect the surface of the ventilator on a daily basis. Treat disposable catheters as infectious medical waste, and send reusable catheters to the disinfection center for sterilization after immersion for 30 minutes in a disinfectant containing 1,000–2,000 mg/L of chlorine-containing disinfectant.

- (5) Prone ventilation: Implementation of prone ventilation for more than 12 hours per day can be considered when $\text{PaO}_2/\text{FiO}_2$ ratio is continuously lower than 150 mmHg (1 mmHg = 0.133 kPa) [8]. (i) Prior to implementing prone ventilation, oropharyngeal and endotracheal secretions should be removed first to ensure an unobstructed artificial airway. (ii) Jejunal tube feeding is recommended for patients undergoing prone ventilation. If gastric tube feeding is employed, aspirate the patient's stomach contents before prone ventilation. When the patient is in the prone position, perform feeding in small amounts. (iii) Regularly evaluate and change the patient's prone position from side to side to avoid applying continuous pressure to the same area. Decompression dressings can be applied to pressured areas and the patient's apophysis area to reduce potential stressful skin damage.
- (6) Prevention of ventilator-associated pneumonia (VAP): Keep the head of the bed elevated by 30 to 45° [13]. Routinely monitor and maintain the cuff pressure of the endotracheal tube at 25–30 cmH₂O. Promptly clear secretions accumulated under the glottis [14].

3.4.2. Maintaining a clear airway

- (1) Cough and expectoration: Teach patients the correct methods of deep breathing and effective coughing based on their condition. In particular, a sealed plastic bag should be used when patients expectorate to prevent the sputum from spreading the virus.
- (2) Atomization inhalation: Administration of drugs via atomization inhaler may produce aerosols and pollute indoor air. Therefore, atomization inhalation should be avoided. If unavoidable, a quantitative inhalation device combined with an aerochamber is preferred.
- (3) Sputum removal: Use a closed endotracheal suctioning system to aspirate secretions in the airway, and perform the procedure only when necessary.

3.4.3. Fever nursing

- (1) Patients with fever should be provided with antipyretic treatment as per medical orders and nursing assessment findings.
- (2) Following antipyretic treatment, closely monitor patients' body temperature, sweating, and electrolyte changes. Dry patients' sweat and change clothes and bed sheets upon heavy sweating. Use gentle actions when replacing clothes and bedding. Treat contaminated clothes and sheets as infectious fabrics and place them into orange waste bags [15].
- (3) Encourage patients who are capable of independent oral food intake to drink more water, and report to the doctor for timely fluid supplements if necessary.

3.4.4. Special care

3.4.4.1. Nursing care for patients with blood purification treatment

- (1) Central venous catheterization should be performed by two nurses, strictly implementing maximum aseptic barrier techniques and protective measures [16].
- (2) Prior to treatment, monitor patients' vital signs, biochemical indicators, and coagulation functions, and evaluate the patency of vascular access.
- (3) Choose the most appropriate anticoagulation approach, treatment mode, and treatment dose as per medical advice.
- (4) During treatment, closely monitor patients' vital signs, coagulation functions, electrolyte changes, treatment effects, and so on. Promptly identify and deal with treatment-related complications such as hypotension, electrolyte disturbance, and thrombocytopenia [17].
- (5) Treatment devices should be used exclusively for the same disease. After treatment, wipe and disinfect the instrument with a disinfectant containing 1,000-2,000 mg/L of chlorine, and discard waste liquid bags and filters in double-layer yellow medical waste bags.

3.4.4.2. Nursing care for patients with convalescent plasma therapy

- (1) Prior to transfusion of convalescent plasma, ensure that patients sign an informed consent form and administer anti-allergic drugs as per medical orders.
- (2) Strictly follow aseptic operation procedures and blood transfusion technical specifications.
- (3) During treatment, particularly within the first 15-20 minutes of the transfusion, closely monitor patients' vital signs, consciousness, and skin changes. Transfuse convalescent plasma slowly within 1-4 hours [17].
- (4) After treatment is completed, discard blood bags in double-layer yellow medical waste bags instead of recycling them. Keep careful records to ensure that donor and recipient information is traceable [17].
- (5) Closely monitor patients' condition after treatment to evaluate the effectiveness of the intervention.

3.4.4.3. Nursing care for patients with extracorporeal membrane oxygenation

Venous-venous extracorporeal membrane oxygenation is the preferred method of respiratory support for patients with severe COVID-19[8].

- (1) Place the catheter carefully to avoid pulling, shifting, bending, leaking, or dropping. Check catheter connections regularly to ensure tightness.
- (2) Closely observe patients' consciousness, pupils, breathing, blood pressure, body temperature, blood oxygen saturation, central venous pressure, mean arterial pressure, and other vital signs, and monitor arterial blood gas analysis results, coagulation functions, and so on [18].
- (3) Device monitoring during extracorporeal membrane oxygenation: (i) Closely monitor the speed of both the centrifugal pump head and the blood flow. (ii) Monitor the inlet and outlet pressure of the oxygenator to identify any obstruction in the device, thereby preventing bending and poor perfusion. (iii) Set the air flow rate and oxygen concentration into the oxygenator based on the incoming blood

flow. (iv) Closely monitor and adjust the temperature of the blood warmer. (v) Prevent and detect complications in a timely manner.

3.4.5. Observation and nursing for drug administration

- (1) Administer drugs according to medical orders, and strictly follow the check system.
- (2) Pharmacy Intravenous Admixture Services (PIVAS) is recommended to ensure accurate dosage and reduce unnecessary exposure.
- (3) Pay attention to drug contraindications. Do not use antiviral drugs such as lopinavir and ritonavir in combination with drugs such as alfuzosin [8]. When traditional Chinese medicine injection is used in conjunction, flush the catheter with normal saline. Separate administration of intestinal microecological regulators by 2 hours from administration of antibiotics.
- (4) Closely monitor adverse drug reactions. When administering antiviral and antibacterial drugs, regularly monitor patients' gastrointestinal responses and liver and kidney functions. When administering glucocorticoids, provide calcium and vitamin D supplements and regularly monitor patients' blood calcium concentration [20].
- (5) Traditional Chinese medicine injections should go through infusion sets with a precision filter. Perform the first 30 minutes of the injection via slow-drip infusion. Do not prescribe traditional Chinese medicine to pregnant women, children, and people with drug allergies.
- (6) One dose of Chinese herbal medicine is allowed per day, delivered separately in 2-4 instances by either warm oral intake or nasal feeding. During drug administration, a light diet is recommended; a cold and greasy diet should be avoided.
- (7) Administer intestinal microecological regulators with warm water not exceeding 40° C. Patients should be helped or instructed when taking the medication.
- (8) Observe the therapeutic effect of administered drugs by closely monitoring patients' body temperature, blood oxygen saturation, and improvement of uncomfortable symptoms.

3.4.6. Management of catheters

- (1) Assess necessity of catheter retention on a daily basis.
- (2) Ensure that all catheters are unobstructed, tightly connected, and fixed firmly to avoid shifting and dropping.
- (3) Closely monitor the color, property, and quantity of the drainage fluid.
- (4) Strictly follow operation and disinfection regulations to avoid secondary infections.
- (5) Use disposable drainage bags and bottles. Strictly implement personal protection measures, use gentle movements, and follow standard operations during catheter replacement to avoid contamination.
- (6) Discard contaminated drainage bags and bottles in double-layer yellow medical waste bags and dispose of them in accordance with relevant regulations.

3.4.7. Observation of disease condition

- (1) Closely monitor changes in patients' vital signs and consciousness, especially breathing frequency, rhythm, pattern, and depth.
- (2) Observe the therapeutic effect of oxygen therapy and dynamically adjust the mode and oxygen flow rate based on patients' degree of dyspnea, blood oxygen saturation, and blood gas analysis results.
- (3) Perform hemodynamic monitoring to detect arrhythmia and heart failure [5].
- (4) Observe associated symptoms such as general muscle pain, fatigue, cough, and chest tightness.
- (5) Record patients' intake and output amounts within 24 hours to maintain water, electrolyte, and acid-base balances.
- (6) Strengthen monitoring of infection indicators and collect blood, sputum, urine, stool, and other specimens promptly and correctly.
- (7) Strengthen monitoring and nursing care of basic diseases, such as hypertension, diabetes, coronary heart disease, and so on.
- (8) Seek to prevent and promptly identify complications. In particular, be vigilant of disease exacerbation when patients develop symptoms such as persistent high fever, respiratory failure, shock, or multiple organs failure [22].

3.4.8. Diet and nutrition

3.4.8.1. For patients with oral food intake

- (1) Ensure patients' oxygen supply is not interrupted when eating or drinking.
- (2) Patients should have small frequent meals, with a diet predominantly composed of egg, beancurd, dairy products, fruit juice, vegetable juice, rice noodles, and so on.
- (3) If nutritional demands are not met, oral or enteral nutrition can be administered.

3.4.8.2. Enteral nutrition support

- (1) Enteral nutrition can be prescribed if patients are normally unable to eat or cannot orally intake at least 60% of their energy consumption for 3-5 consecutive days.
- (2) Monitor nutrition indicators, including albumin, transferrin, and preprotein counts. Record the type, concentration, and quantity of enteral nutrition supplement [24].
- (3) During enteral nutrition infusion, elevate the head of bed at an angle $\geq 30^\circ$ to prevent accidental aspiration.
- (4) Observe development of gastrointestinal symptoms such as diarrhea, nausea, or vomiting.
- (5) A permissive low-calorie program that can satisfy 60%–80% of nutritional intake is recommended during the early stage. Once the disease is alleviated, energy and nutrients can be gradually increased to the full amount.

3.4.8.3. Parenteral nutrition support

- (1) Parenteral nutrition is required for patients with severe gastrointestinal dysfunction.
- (2) Closely monitor any development of metabolic disorders and complications.

3.4.9. Basic nursing

3.4.9.1. Bed allocation

Single rooms are recommended. However, if single rooms are not available, bed spacing should be at least 1 m. A negative pressure ward is recommended, allowing at least 12 air exchange cycle per hour. For non-negative pressure wards, air flow rate must remain at least 160 L/s during natural air ventilation [25].

3.4.9.2. Oral care

- (1) Patients who can do self-care for themselves should be encouraged to perform oral care themselves.
- (2) Patients who cannot care for themselves should be assisted by nurses to perform oral care 2 or 3 times per day.
- (3) Be vigilant of potential splash of secretions when performing oral care for patients. Nurses should wear Level III personal protection equipment if necessary.

3.4.9.3. Skin care

- (1) Closely monitor facial skin for any pressure injury during oxygen therapy.
- (2) Observe patients' overall skin condition and instruct them to either change position regularly or use protective products to prevent skin damage.

3.4.9.4. Elimination care

- (1) Prevent constipation. During and after defecation, carefully increase patients' oxygen concentration to prevent hypoxia.
- (2) Prevent diarrhea. If patients develop diarrhea, administer drugs as per medical orders and be vigilant to prevent incontinence dermatitis.
- (3) Bedpans should be used exclusively for each patient. After use, immerse the bedpan in a disinfectant solution containing 5,000 mg/L of effective chlorine and subsequently clean it for future use.
- (4) If the medical institution is equipped with a sewage treatment system, excreta can be discharged directly into the sewage tank. If not, excreta must be disinfected in accordance with the *Regulation of Disinfection Technique in Healthcare Settings* [26].

3.4.10. Psychological nursing

- (1) Introduce patients to the ward environment and isolation regulations upon admission.
- (2) Regularly assess patients' psychological condition and consult with the psychiatry department if necessary.
- (3) During bedside operations, provide patients with appropriate emotional support through gestures such as eye contact, touch, nods, handshakes, and thumbs-up gestures, thereby helping them to build confidence in their ability to overcome the disease [27].
- (4) Help to communicate information to external relatives, provide patients with continuous information support, and encourage them to actively collaborate in their treatment.

3.4.11. Respiratory rehabilitation

- (1) Principle: Collaborate with the clinical medical team to assess the feasibility of commencing respiratory rehabilitation program for patients. Respiratory rehabilitation activities should not reduce the patient's blood oxygen saturation and blood pressure [28].
- (2) Instructional format: Videos, brochures, etc.
- (3) Rehabilitation contents: Routine changes in body posturing (e.g., semi-recumbent position, lateral position, prone position, etc.) to reduce the work of breathing muscles and save energy; respiratory control techniques (e.g., abdominal breathing and pursed-lip breathing) to expand the lower chest and relieve breathing difficulties; effective sputum excretion techniques (e.g., postural drainage and effective coughing) to promote sputum excretion and maintain an unobstructed airway.

3.4.12. Rest and activities

- (1) To ensure adequate sleep, instruct patients to rest in bed. Those with sleep disorders should be managed by drugs as per medical orders.
- (2) Patients who can get out of the bed should follow the "Three Steps to Get Up" guideline (lie in the bed for 30 seconds before getting up, drop both legs for 30 seconds before standing up, and stand for 30 seconds before walking). Instruct patients to perform activities such as sitting, standing, and stepping along the bedside once they are out of bed [29].
- (3) Prevent patients from falling out of bed. Instruct patients to perform muscle training such as clenching their fists, raising their arms, ankle pumping, heel slipping, lifting their legs, and contracting quadriceps and gluteal muscles, according to their tolerance levels. Compression stockings can be utilized to prevent the formation of deep vein thrombosis in lower limbs.
- (4) Avoid excessive activities to reduce patients' blood oxygen saturation and blood pressure.

3.4.13 Discharge instructions

- (1) Instruct patients to perform respiratory rehabilitation activities according to their rehabilitation program.
- (2) Ensure timely transmission of patients' medical information and instruct them to go to the designated place for centralized isolation of 14 days [30].
- (3) Inform patients of follow-up instructions for weeks 2 and 4 and carefully perform follow-up and review.
- (4) Instruct patients to work and rest regularly and ensure they maintain adequate sleep, a balanced diet, and a calm emotional state.
- (5) Instruct patients to be vigilant of hand hygiene, maintain distance from family members, wear a mask, have separate meals, and avoid sharing personal items such as tableware and washing materials during home isolation [31].

Expert Consensus Panel on Holistic Care of COVID-19 Patients in Severe and Critical Condition (in alphabetical order of surnames)

Miaoxia Chen (the Third Affiliated Hospital of Sun Yat-sen University), Man Cui (Peking University Third Hospital), Lu Deng (the Second Xiangya Hospital of Central South University), Xin Fu (China-Japan Union Hospital of Jilin University), Lan Gao (the First Hospital of Jilin University), Qiuying Han (Zhongshan Hospital, Xiamen University), Daling Hu (Sir Run Run Hospital, Nanjing Medical University), Guimei Li (the People's No. 3 Hospital of Anyang), Minmin Li (Shandong Provincial Hospital), Zhengyin Liu (Peking Union Medical College Hospital, the Chinese Academy of Medical Sciences), Hong Sun (Peking Union Medical College Hospital, the Chinese Academy of Medical Sciences), Xinjuan Wu (Peking Union Medical College Hospital, the Chinese Academy of Medical Sciences), Hui Wang (Tongji Hospital Affiliated with Tongji Medical College of Huazhong University of Science and Technology), Ling Wang (Peking University People's Hospital), Yuying Wang (Peking University First Hospital), Xia Wang (Beijing Hospital), Yanjun Wang (the Second Hospital of Jilin University), Lili Wei (the Affiliated Hospital of Qingdao University), Huiyun Yang (the Second Affiliated Hospital of Xi'an Jiaotong University), Tieying Zeng (Tongji Hospital Affiliated with Tongji Medical College of Huazhong University of Science and Technology), Yinghui Zhang (the Second Hospital of Shanxi Medical University), Jianping Zhao (Tongji Hospital Affiliated with Tongji Medical College of Huazhong University of Science and Technology), Peiyu Zhao (China-Japan Friendship Hospital), Zhuqing Zhong (the Third Xiangya Hospital of Central South University).

References

- [1] World Health Organization. Novel coronavirus (2019-nCoV) situation report - 12 [Internet]. Available from: https://www.who.int/docs/default-source/coronaviruse/situation-reports/2020-0201-sitrep-12-ncov.pdf?sfvrsn=273c5d35_2.
- [2] National Health Commission of the People's Republic of China. Announcement of the National Health Commission of the People's Republic of China, No. 1, 2020 [Internet]. Available from: <http://www.nhc.gov.cn/jkj/s7916/202001/44a3b8245e8049d2837a4f27529cd386.shtml>.
- [3] Chinese Preventive Medicine Association, Expert Group on the Prevention and Control of Novel Coronavirus Pneumonia. Latest understanding of epidemiological characteristics of novel coronavirus pneumonia. Chin J Vir Dis. doi: 10.16505/j.2095-0136.2020.0015. Epub 2020 Feb 24.

- [4] Yang X, Yu Y, Xu J, Shu H, Xia J, Liu H, et al. Clinical course and outcomes of critically ill patients with SARS-CoV-2 pneumonia in Wuhan, China: a single-centered, retrospective, observational study. *Lancet Respir Med*. 2020 Feb 24. doi: 10.1016/S2213-2600(20)30079-5.
- [5] National Health Commission of the People's Republic of China. Diagnosis and Treatment Protocol for Novel Coronavirus Pneumonia (Trial version 6) [Internet]. Available from: <http://www.nhc.gov.cn/xcs/zhengcwj/202002/8334a8326dd94d329df351d7da8aefc2/files/b218cfcb1bc54639af227f922bf6b817.pdf>.
- [6] Tongji Hospital Affiliated with Tongji Medical College of Huazhong University of Science and Technology, Novel Coronavirus Pneumonia Treatment Collaboration Group. Consensus on the diagnosis and management of severe novel coronavirus pneumonia cases [Internet]. Available from: <http://health.people.com.cn/n1/2020/0211/c14739-31581678.html>.
- [7] Jin Y, Cai L, Cheng Z, Cheng H, Deng T, Fan Y, et al. Quick guide to the diagnosis and treatment of novel coronavirus (2019-nCoV) pneumonia (standard version). *Med J Chin People Lib Army*. Epub 2020 Feb 2. Available from: <http://kns.cnki.net/kcms/detail/11.1056.r.20200201.1338.003.html>.
- [8] National Health Commission of the People's Republic of China. Diagnosis and treatment schemes for severe novel coronavirus-infected pneumonia cases (trial) [Internet]. Available from: <http://www.nhc.gov.cn/yzygj/s7653p/202001/9fbefc9a5fe747e98ea5baeedfb68158/files/4967f0d3f24646819c132a7e1797440d.pdf>.
- [9] Respiratory Critical Care Medicine Group of the Respiratory Medicine Branch of the Chinese Medical Association, Critical Care Medicine Working Committee of the Respiratory Medicine Branch of the Chinese Medical Doctor Association. Airway management recommendations for adults with severe novel coronavirus pneumonia (trial). *Natl Med J China* 2020;100:E004.
- [10] Ni Z, Qin H, Li J, Wang Q, Wang J, Jing G, et al. Expert consensus on the management of nasal high-flow oxygen therapy in patients with novel coronavirus pneumonia. *Chin J Respir Crit Care Med* 2020;19:110-5.
- [11] Ge H, Dai B, Xu P, Duan K, Xia J, Duan J, et al. Expert consensus on the sensory management of ventilator use in patients with novel coronavirus pneumonia. *Chin J Respir Crit Care Med*. 2020;19:116-9.
- [12] Wang Q, Jing X, Zhu Z, Zheng J, Kong R, Zhang X, et al. Discussions on key infection prevention and control issues of ventilator use in novel coronavirus pneumonia patients under mechanical ventilation. *Chin J Infect Dis*. 2020;38:E005.
- [13] Infectious Diseases Group of the Respiratory Medicine Branch of the Chinese Medical Association. Diagnosis and treatment guidelines for hospital-acquired pneumonia and ventilator-related pneumonia in Chinese adults (2018 edition). *Chin J Tuberc Respir Dis*. 2018;41(4):255-80.

- [14] Ni Z, Luo F, Wang J, Liu T, Zhang T, Yang W, et al. Suggestions on aerosol inhalation therapy for patients with novel coronavirus infection. *Chin J Respir Crit Care Med*. 2020;19:120-4.
- [15] Regulation for washing and disinfection technique of medical textiles in healthcare facilities (WS/T 508-2016). *Chin J Infect Control* 2017;16(7):687-92.
- [16] American Society of Anesthesiologists. Practice guidelines for central venous access 2020: an updated report by the American Society of Anesthesiologists task force on central venous access. *Anesthesiology* 2020;132:8-43.
- [17] Expert Team of Chinese Medical Association Nephrology Branch. [Recommendations for prevention and control of novel coronavirus infection in blood purification center (room) from the Chinese Medical Association Nephrology Branch]. *Chin J Nephrol* 2020;36:82-4.
- [18] Zhang W, Pan C, Song Q. Key issues during respiratory therapy for patients with novel coronavirus pneumonia. *Med J Chin People Lib Army*. Epub 2020 Feb 27. Available from: <http://kns.cnki.net/kcms/detail/11.1056.r.20200226.1950.004.html>.
- [19] Committee of Critical Care Medicine, Chinese Association of Chest Physician, Chinese Medical Doctor Association, Critical Care Medicine Group, Chinese Thoracic Society, Chinese Medical Association. [Recommendations for clinical application of extracorporeal membrane oxygenation in adults severe respiratory failure]. *Chin J Tuberc Respir Dis*. 2019;42:660-84.
- [20] Meng R, Wang J. Progress in the prevention and treatment of glucocorticoid-induced osteoporosis. *Chin J Osteoporos*. 2019;25:123-6.
- [21] Chinese Pharmacopoeia Commission. Pharmacopoeia of the People's Republic of China, vol. 1. Beijing: Chinese Medical Science and Technology Press; 2015.
- [22] Ma J, Hu F, Sun HM, Chen J, Ding XB, Li J. Nursing experience of treating a critical ill patient with novel coronavirus (2019-nCoV) infected pneumonia by using ECMO. *J New Med* 2020;30:74-7.
- [23] Xu J, Zeng F, Wu Y, Zhou T, Han Y, Lu Y, et al. Nutritional Support and Monitoring Recommendations for Patients with Severe Novel Coronavirus-infected Pneumonia. *Chin J Hosp Pharm*. Epub 2020 Feb 19. Available from: <http://kns.cnki.net/kcms/detail/42.1204.r.20200218.0954.002.html>.
- [24] Chen W. Clinical application of nutritional dietary guidelines for the prevention and treatment of novel coronavirus-infected pneumonia. *Beijing Med J*. Epub 2020 Feb 19. doi: 10.15932/j.0253-9713.2020.03.040.
- [25] World Health Organization. Infection prevention and control during health care when novel coronavirus (nCoV) infection is suspected. Available from: [https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-\(ncov\)-infection-is-suspected-20200125](https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-(ncov)-infection-is-suspected-20200125)

- [26] National Health Commission of the People's Republic of China. Regulation of Disinfection Technique in Healthcare Settings [Internet]. Available from: <http://www.nhc.gov.cn/xxgk/pages/viewdocument.jsp?dispatchDate=&staticUrl=/zwgkzt/s9496/201204/54510.shtml>.
- [27] National Health Commission of the People's Republic of China. Notice on issuing guiding principles for emergency psychological crisis interventions for patients with novel coronavirus-infected pneumonia [Internet]. Available from: http://www.gov.cn/xinwen/2020-01/27/content_5472433.htm.
- [28] Wang C, Fang F, Xie Y, Zhao H, Yu P. Guidelines for respiratory rehabilitation of patients with 2019 novel coronavirus pneumonia (first edition). *Chin J Repar Reconstr Surg* 2020;34(3):275-9.
- [29] Tan H, Chen Y, Chen S, Mo H, Li X, Tang X. Development and application of fall prevention process management among hospitalized patients. *Chin Nurs Manag.* 2017;17:818-21.
- [30] National Health Commission of the People's Republic of China. Performing high quality follow-up of discharged patients with novel coronavirus pneumonia [Internet]. Available from: http://www.gov.cn/zhengce/zhengceku/2020-02/20/content_5481166.htm.
- [31] Chinese Health Education Center, National Health Commission Ministry of Publicity. Health education manual for novel coronavirus-infected pneumonia [Internet]. Available from: http://www.gov.cn/fuwu/2020-02/10/content_5476794.htm.